

Retirement age and type as predictors of frailty : a retrospective cohort study of older businessmen

Haapanen, Markus J.

2020

Haapanen , M J , von Bonsdorff , M B , Perttilä , N M , Törmäkangas , T , von Bonsdorff , M E , Strandberg , A Y & Strandberg , T E 2020 , ' Retirement age and type as predictors of frailty : a retrospective cohort study of older businessmen ' , BMJ Open , vol. 10 , no. 12 , 037722 . <https://doi.org/10.1136/bmjopen-2020-037722>

<http://hdl.handle.net/10138/326039>

<https://doi.org/10.1136/bmjopen-2020-037722>

cc_by_nc

publishedVersion



Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

BMJ Open Retirement age and type as predictors of frailty: a retrospective cohort study of older businessmen

Markus J Haapanen ^{1,2}, Mikaela B von Bonsdorff,^{2,3} Niko M Perttälä,¹ Timo Törmäkangas,³ Monika E von Bonsdorff,^{4,5} Arto Y Strandberg ⁶, Timo E Strandberg^{6,7}

To cite: Haapanen MJ, von Bonsdorff MB, Perttälä NM, et al. Retirement age and type as predictors of frailty: a retrospective cohort study of older businessmen. *BMJ Open* 2020;10:e037722. doi:10.1136/bmjopen-2020-037722

Prepublication history and supplemental material for this paper is available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2020-037722>).

Received 13 February 2020
Revised 16 November 2020
Accepted 27 November 2020



* Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Markus J Haapanen;
markus.haapanen@helsinki.fi

ABSTRACT

Objectives To study the association between retirement characteristics and frailty in a homogenous population of former business executives.

Design Cross-sectional cohort study using data from the Helsinki Businessmen Study.

Setting Helsinki, Finland.

Participants 1324 Caucasian men, born in 1919–1934, who had worked as business executives and managers and of whom 95.9% had retired by the year 2000.

Questions on age at and type of retirement, lifestyle and chronic conditions were embedded in questionnaires.

Primary and secondary outcome measures Frailty assessed according to a modified phenotype definition at mean age 73.3 years.

Results Mean age at retirement was 61.3 years (SD 4.3) and 37.1% had retired due to old age. The prevalence of frailty was lowest among men retiring at ages 66–67 years but increased among those who worked up to age 70 years or older. Compared with men who retired before age 55 years, those retiring at ages 58–69 years were at decreased risk of frailty in old age relative to non-frailty (adjusted ORs 0.07–0.29, $p < 0.05$). Compared with men who transitioned into old age retirement, those who retired due to disability were at increased risk of prefrailty (adjusted OR 1.53, 95% CI 1.01 to 2.32) and frailty (adjusted OR 3.52, 95% CI 1.97 to 6.29), relative to non-frailty.

Conclusion Exiting working life early and continuing to be occupationally active until age 70 years and older were both associated with increased risk of frailty among the men. Promotion of longer work careers could, however, promote healthier ageing, as the lowest prevalence of frailty was observed in former business executives who retired at ages 66–67 years.

INTRODUCTION

Strengths and limitations of this study

This study is among the first to report associations between the age at retirement, type of pension and frailty in old age among a homogenous population characterised with high socioeconomic position.

Generalisation of the results to other socioeconomic or occupational groups should be made with caution. The study included 1324 men who had worked as business executives and managers during their work careers.

Frailty was defined according to a modified phenotype definition, assessed using questionnaire data, and information on retirement characteristics was self-reported in this cross-sectional study.




Figure 1 Distribution of frailty classification assessed at an average age of 73.3 years according to categories of age at retirement.

METHODS

Study population

Age at and type of retirement

Frailty




Table 1 Characteristics of the men according to retirement type (proportion and per cent unless stated otherwise)

n (%)	All	Old age retirement	Early old age retirement	Pension bene t package	Unemployment retirement	Disability retirement	P value*
Old age characteristics assessed in the year 2000 (n=1324)							
Age in years, mean (SD)	73.3 (4.1)	74.6 (4.0)	72.9 (4.0)	72.5 (3.9)	71.4 (3.1)	72.6 (4.1)	<0.001
Cardiovascular disease, n (%)	812 (59.4)	307 (60.6)	159 (55.2)	155 (61.8)	47 (45.6)	124 (70.9)	<0.001
Diabetes, n (%)	113 (8.6)	35 (6.9)	22 (7.6)	26 (10.4)	7 (6.8)	23 (13.1)	0.018
Smoking status, n (%) (n=1322)							
Current smoker	106 (8.0)	35 (6.9)	19 (6.6)	27 (10.8)	10 (9.7)	15 (8.6)	0.371
Quit smoking	719 (54.8)	269 (53.3)	160 (55.6)	129 (51.4)	59 (57.3)	102 (58.3)	
Never smoked	497 (37.2)	201 (39.8)	109 (37.8)	95 (37.8)	34 (33.0)	58 (33.1)	
Alcohol consumption, n (%) (n=1040)							
None	153 (14.7)	68 (16.9)	30 (12.9)	23 (11.3)	11 (13.1)	21 (17.8)	<0.001
Light	448 (43.1)	176 (43.8)	114 (48.9)	79 (38.9)	36 (42.9)	43 (36.4)	
Moderate	234 (22.5)	103 (25.6)	39 (16.7)	47 (23.2)	18 (21.4)	27 (22.9)	
High	205 (19.7)	55 (13.7)	50 (21.5)	54 (26.6)	19 (22.6)	27 (22.9)	
Age at retirement (n=1284)							
In years, mean (SD)	61.3 (4.3)	63.5 (3.6)	61.0 (3.7)	60.6 (3.7)	59.1 (3.7)	57.7 (4.4)	<0.001
Categorised age at retirement, n (%) (n=1284)							
70 years	23 (1.8)	14 (2.9)	4 (1.4)	5 (2.0)	0	0	<0.001
68-69 years	15 (1.2)	11 (2.3)	1 (0.4)	2 (0.8)	0	1 (0.6)	
66-67 years	52 (4.1)	40 (8.2)	5 (1.8)	7 (2.9)	0	0	
64-65 years	273 (21.3)	204 (41.8)	35 (12.5)	21 (8.6)	6 (6.0)	7 (4.1)	
62-63 years	292 (22.8)	126 (25.8)	82 (29.2)	45 (18.4)	15 (15.0)	24 (14.0)	
60-61 years	321 (24.9)	69 (14.1)	85 (30.2)	90 (36.9)	38 (38.0)	39 (22.8)	
58-59 years	127 (9.9)	11 (2.3)	41 (14.6)	37 (15.2)	13 (13.0)	25 (14.6)	
56-57 years	87 (6.8)	3 (0.6)	17 (6.0)	20 (8.2)	17 (17.0)	30 (17.5)	
55 years	94 (7.3)	10 (2.0)	11 (3.9)	17 (7.0)	11 (11.0)	45 (26.3)	
Frailty assessed in the year 2000, n (%) (n=1324)							
Non-frail	496 (37.5)	177 (34.9)	108 (37.5)	123 (49.0)	44 (42.7)	44 (25.1)	<0.001
Prefrail	686 (51.8)	281 (55.4)	152 (52.8)	110 (43.8)	49 (47.6)	94 (53.7)	
Frail	142 (10.7)	49 (9.7)	28 (9.7)	18 (7.2)	10 (9.7)	37 (21.1)	
Stressfulness of work career, assessed in the year 2003 (n=972)							
0 (high stress) to 10 (no stress)	5.3 (2.8)	5.7 (2.9)	5.2 (2.8)	4.9 (2.7)	5.7 (3.0)	4.6 (2.7)	<0.001

*P for difference in means/distributions between groups.

Table 2 ORs and 95% CIs of prefrailty and frailty compared with non-frailty according to age at and type of retirement

	Prefrailty		Frailty	
	OR (95% CI)		OR (95% CI)	
	Model 1	Model 2	Model 1	Model 2
Categories of retirement age (years)				
70	0.78 (0.25 to 2.46)	0.83 (0.26 to 2.62)	0.28 (0.06 to 1.40)	0.38 (0.08 to 1.97)
68 69	0.51 (0.15 to 1.73)	0.55 (0.16 to 1.89)	0.11 (0.01 to 1.03)	0.10 (0.01 to 1.02)
66 67	0.49 (0.23 to 1.04)	0.52 (0.24 to 1.12)	0.07 (0.01 to 0.34)***	0.07 (0.01 to 0.37)**
64 65	0.37 (0.21 to 0.64)***	0.40 (0.22 to 0.70)***	0.19 (0.09 to 0.40)***	0.21 (0.10 to 0.46)***
62 63	0.61 (0.35 to 1.06)	0.67 (0.38 to 1.17)	0.20 (0.09 to 0.45)***	0.24 (0.11 to 0.54)***
60 61	0.49 (0.29 to 0.85)**	0.48 (0.28 to 0.83)**	0.22 (0.10 to 0.47)***	0.22 (0.10 to 0.47)***
58 59	0.45 (0.24 to 0.83)**	0.46 (0.25 to 0.87)*	0.28 (0.12 to 0.66)**	0.29 (0.12 to 0.70)**
56 57	0.50 (0.26 to 0.98)*	0.49 (0.25 to 0.98)*	0.51 (0.21 to 1.24)	0.45 (0.18 to 1.14)
55	(Ref)	(Ref)	(Ref)	(Ref)
Type of retirement				
Old age	(Ref)	(Ref)	(Ref)	(Ref)
Early old age	1.02 (0.74 to 1.40)	0.99 (0.72 to 1.37)	1.22 (0.71 to 2.09)	1.24 (0.72 to 2.16)
Pension benefit package	0.66 (0.47 to 0.91)*	0.61 (0.44 to 0.85)**	0.71 (0.39 to 1.30)	0.62 (0.33 to 1.15)
Unemployment	0.89 (0.56 to 1.41)	0.86 (0.54 to 1.37)	1.36 (0.62 to 2.98)	1.44 (0.64 to 3.21)
Disability	1.62 (1.07 to 2.45)*	1.46 (0.96 to 2.23)	4.34 (2.47 to 7.63)***	3.52 (1.97 to 6.29)***
Stressfulness of work career				
Per 1-unit increase	0.94 (0.89 to 0.99)*	0.94 (0.90 to 0.99)*	0.94 (0.86 to 1.03)	0.95 (0.87 to 1.04)

*P<0.001; **P<0.01; ***P<0.05.

Model 1 adjusted for age in the year 2000 (age at measuring frailty).

Model 2 adjusted additionally for smoking, alcohol consumption, cardiovascular disease and diabetes.

Age at retirement and frailty in old age

Type of retirement and frailty in old age




Figure 2 Distribution of frailty classification assessed at an average age of 73.3 years according to the type of retirement.

Stressfulness of work career

DISCUSSION

Author affiliations

¹Department of General Practice and Primary Health Care, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

²Folkhälsan Research Centre, Helsinki, Finland

³Department of Health Sciences, Gerontology Research Centre, University of Jyväskylä Faculty of Sport and Health Sciences, Jyväskylä, Finland

⁴School of Business and Kokkola University Consortium Chydenius, University of Vaasa, Vaasa, Finland

⁵Department of Management and Leadership, Jyväskylä University School of Business and Economics, Jyväskylä, Finland

⁶Department of Medicine, University of Helsinki and Helsinki University Hospital, Helsinki, Finland

⁷Center for Life Course Health Research, University of Oulu, Oulu, Finland

Twitter Mikaela B von Bonsdorff @vonbonsdorffx2

Contributors MJH designed the present study, drafted the article, analysed the data and drafted the manuscript. MBvB designed the present study, interpreted the data and revised the article critically for important intellectual content. MEvB, NMP, TT and AYS interpreted the data and revised the article critically for important intellectual content. TES acquired funding for the cohort, data collection, interpreted the data and revised the article critically for important intellectual content.

Funding The Academy of Finland supported MBvB with grant 257239. The Academy of Finland supported TT with grant 286536. The Academy of Finland supported MEvB with grants 294530, 307114 and 303920. TES is supported by the Helsinki University Hospital, Department of Medicine.

Disclaimer The sponsor had no role in the design, methods, subject recruitment, data collections, analysis and preparation of the paper.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval The Ethics Committee of the Department of Medicine, University of Helsinki, has approved the research protocol of the follow-up study of HBS. All participants provided a written informed consent before participating in the study. The study adheres with the principles stated in the Declaration of Helsinki.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Data may be obtained from a third party and are not publicly available. Inquiries regarding the data sets used and/or analysed during the current study can be directed to the principal investigator (TES) of the Helsinki Businessmen Study.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iDs

Markus J Haapanen <http://orcid.org/0000-0002-8632-7906>

Arto Y Strandberg <http://orcid.org/0000-0002-6080-1729>

REFERENCES

- Fried LP, Tangen CM, Walston J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001;56:M146-57.
- Bergman H, Ferrucci L, Guralnik J, et al. Frailty: an emerging research and clinical paradigm-issues and controversies. *J Gerontol A Biol Sci Med Sci* 2007;62:731-7.
- United Nations, Department of Economic and Social Affairs PD. World population prospects: the 2015 revision, key findings and advance tables Working Paper No. ESA/P/WP.241. New York, 2015.
- Morley JE, Vellas B, van Kan GA, et al. Frailty consensus: a call to action. *J Am Med Dir Assoc* 2013;14:392-7.
- Cesari M, Prince M, Thiagarajan JA, et al. Frailty: an emerging public health priority. *J Am Med Dir Assoc* 2016;17:188-92.
- Oshio T, Kan M. The dynamic impact of retirement on health: evidence from a nationwide ten-year panel survey in Japan. *Prev Med* 2017;100:287-93.
- Rechel B, Grundy E, Robine J-M, et al. Ageing in the European Union. *Lancet* 2013;381:1312-22.
- Cesari M, Prince M, Thiagarajan JA, et al. Frailty: an emerging public health priority. *J Am Med Dir Assoc* 2016;17:188-92.
- Schinkel-Ivy A, Mosca I, Mansfield A. Factors contributing to unexpected retirement and unemployment in adults over 50 years old in Ireland. *Gerontol Geriatr Med* 2017;3:2333721417722709:233372141772270.
- Lu W, Benson R, Glaser K, et al. Relationship between employment histories and frailty trajectories in later life: evidence from the English longitudinal study of ageing. *J Epidemiol Community Health* 2017;71:439-45.
- Official Statistics of Finland (OSF). Labour Force Survey [e-Publication]. Helsinki, Finland, 2016. http://tilastokeskus.fi/til/tyti/index_en.html
- Virtanen M, Oksanen T, Pentti J, et al. Occupational class and working beyond the retirement age: a cohort study. *Scand J Work Environ Health* 2017;43:426-35.
- Strandberg TE, Salomaa V, Strandberg AY, et al. Cohort profile: the Helsinki businessmen study (HBS). *Int J Epidemiol* 2016;45:1074-1074h.
- Miettinen TA, Huttunen JK, Naukkarinen V, et al. Multifactorial primary prevention of cardiovascular diseases in middle-aged men. risk factor changes, incidence, and mortality. *JAMA* 1985;254:2097-102.
- von Bonsdorff MB, Haapanen MJ, Törmäkangas T, et al. Midlife cardiovascular status and old age physical functioning trajectories in older businessmen. *J Am Geriatr Soc* 2019;67:2490-6.
- Sirala J, Pitkälä KH, Tilvis RS, et al. Definition of frailty in older men according to questionnaire data (RAND-36/SF-36): the Helsinki businessmen study. *J Nutr Health Aging* 2011;15:783-7.
- Järnefelt N. Education and longer working lives: a longitudinal study on Education differences in the late exit from working life of older employees in Finland.
- RAND. The Rand 36-Item health survey the Rand 36-Item health survey. *Health* 1992;2:91-7.
- Strandberg AY, Trygg T, Pitkälä KH, et al. Alcohol consumption in midlife and old age and risk of frailty. *Age Ageing* 2018;47:248-54.
- von Bonsdorff MB, von Bonsdorff ME, Haanpää M, et al. Work-loss years among people diagnosed with diabetes: a reappraisal from a life course perspective. *Acta Diabetol* 2018;55:485-91.
- Working Group on Separate Pension Systems. Separate pension systems report of the Working group. Minist Soc Aff Heal 2019 <http://julkaisut.valtioneuvosto.fi/handle/10024/161385>
- Kannisto J. Eläkelän mittaaminen [Measuring retirement age]. Finnish Cent Pensions 2006:156. <https://www.julkari.fi/bitstream/handle/10024/129385/Elakkeellesiirtymisiannittaminen.pdf?sequence=1&isAllowed=y>
- Salonen J. Statutory pensions for the self-employed: working life, con rmed income and nancing. Cent Pensions, Res Reports, 2015. Available: <https://www.etk.fi/en/julkaisu/statutory-pensions-for-the-self-employed-working-life-con rmed-income-and-nancing/>
- Huohvanainen E, Strandberg AY, Stenholm S, et al. Association of self-rated health in midlife with mortality and old age frailty: a 26-Year follow-up of initially healthy men. *J Gerontol A Biol Sci Med Sci* 2016;71:923-8.
- Stolz E, Mayerl H, Waxenegger A, et al. Impact of socioeconomic position on frailty trajectories in 10 European countries: evidence from the survey of health, ageing and retirement in Europe (2004-2013). *J Epidemiol Community Health* 2017;71:73-80.
- de Wind A, van der Pas S, Blatter BM, et al. A life course perspective on working beyond retirement-results from a longitudinal study in the Netherlands. *BMC Public Health* 2016;16:499.
- van der Zwaan GL, Oude Hengel KM, Sewdas R, et al. The role of personal characteristics, work environment and context in working beyond retirement: a mixed-methods study. *Int Arch Occup Environ Health* 2019;92:535-49.
- Collard RM, Boter H, Schoevers RA, et al. Prevalence of frailty in community-dwelling older persons: a systematic review. *J Am Geriatr Soc* 2012;60:1487-92.
- Di Gessa G, Corna L, Price D, et al. The decision to work after state pension age and how it affects quality of life: evidence from a 6-year English panel study. *Age Ageing* 2018;47:450-7.
- Di Gessa G, Corna LM, Platts LG, et al. Is being in paid work beyond state pension age beneficial for health? Evidence from England using a life-course approach. *J Epidemiol Community Health* 2017;71:431-8.
- Burke RJ. Sources of managerial and professional stress in large organizations. In: Causes, coping and consequences of stress at work. New York: John Wiley and Sons Ltd, 1988: 77-114.
- Kulmala J, von Bonsdorff MB, Stenholm S, et al. Perceived stress symptoms in midlife predict disability in old age: a 28-year prospective cohort study. *J Gerontol A Biol Sci Med Sci* 2013;68:984-91.
- von Bonsdorff MB, von Bonsdorff M, Kulmala J, et al. Job strain in the public sector and hospital in-patient care use in old age: a 28-year prospective follow-up. *Age Ageing* 2014;43:393-9.
- von Bonsdorff MB, Seitsamo J, von Bonsdorff ME, et al. Job strain among blue-collar and white-collar employees as a determinant of

- total mortality: a 28-year population-based follow-up. *BMJ Open* 2012;2:e000860.
- 35 Kalousova L, Mendes de Leon C, de Leon CM. Increase in frailty of older workers and retirees predicted by negative psychosocial working conditions on the job. *Soc Sci Med* 2015;124:275–83.
 - 36 Palmer KT, D'Angelo S, Harris EC, et al. Frailty, prefrailty and employment outcomes in health and employment after injury (Heaf) study. *Occup Environ Med* 2017;74:476–82.
 - 37 Szabó Z, Gnes, Allen J, Stephens C, et al. Is retirement associated with physical health benefits? A longitudinal investigation with older new Zealanders. *Age Ageing* 2019;48:267–72.
 - 38 Mosca I, Barrett A. Increasing the state pension age, the recession and expected retirement ages. *Econ Soc Rev* 2013;44:447–72.
 - 39 McMichael AJ. Standardized mortality ratios and the "healthy worker effect": Scratching beneath the surface. *J Occup Med* 1976;18:165–8.